

Ingrid Tayane Vieira da Silva do Nascimento¹ | Nathália Medeiros Guimarães² | Raquel Soares Martins³
Thiago Anchieta de Melo⁴ | Ilka Márcia Ribeiro de Souza Serra⁵ | Maria Claudene Barros⁶

HEALTH OF CULTIVATED OYSTERS: A LITERATURE REVIEW

SANIDADE DE OSTRAS CULTIVADAS: UMA REVISÃO DE LITERATURA

SALUD DE LAS OSTRAS CULTIVADAS: UNA REVISIÓN DE LA LITERATURA

ABSTRACT

Oysters pose a high risk to public health because, as filter-feeding organisms, they accumulate large quantities of microorganisms present in contaminated water. Thus, we aim in this work, conducting research on the presence of fungi, yeast and bacteria isolated from cultivated oysters, through an extensive literature review in the last ten years. We carried out a bibliographical in February 2023, using the descriptors: "bivalve mollusc", "oyster farming", "oyster farming", "fungi", "yeasts" and "bacteria". The studies selected include articles related to the health of oysters cultivated in Brazil and in the world. We collected a total of 719 articles distributed in the Science Direct (395), Scielo (06), Google Scholar (127) and PubMed (191) databases. Only 20 articles met the criteria for inclusion in the research. The papers surveyed were mostly published in Korea, India, and the Philippines, in the year 2020. The study reveals the scarcity of aspect involving fungi and yeasts isolated from cultivated oysters. The selected authors identified bacteria of the genus *Vibrio* spp., *Escherichia coli* and *Salmonella* sp. in most of their works. Thus, it is necessary to increase studies that can assess the risks that such pathogenic organisms may cause in cultivable oysters and prevent damage to human and animal health.

KEYWORDS

Fungi; bacteria; oyster farming.

RESUMO

Ostras representam um alto risco à saúde pública, pois, como organismos filtradores, acumulam grandes quantidades de microrganismos presentes em águas contaminadas. Assim, objetivamos neste trabalho, realizar pesquisas sobre a presença de fungos, leveduras e bactérias isoladas de ostras cultivadas, através de uma extensa revisão de literatura nos últimos dez anos. Realizamos levantamento bibliográfico em fevereiro de 2023, utilizando os descritores: "molusco bivalve", "criação de ostras", "criação de ostras", "fungos", "leveduras" e "bactérias". Os estudos selecionados incluem artigos relacionados à saúde das ostras cultivadas no Brasil e no mundo. Coletamos um total de 719 artigos distribuídos nas bases de dados Science Direct

(395), Scielo (06), Google Scholar (127) e PubMed (191). Apenas 20 artigos atenderam aos critérios de inclusão na pesquisa. Os artigos pesquisados foram publicados em sua maioria na Coreia, Índia e Filipinas, no ano de 2020. O estudo revela a escassez de aspectos envolvendo fungos e leveduras isolados de ostras cultivadas. Os autores selecionados identificaram bactérias do gênero *Vibrio* spp., *Escherichia coli* e *Salmonella* sp. na maioria de suas obras. Assim, faz-se necessário incrementar estudos que possam avaliar os riscos que tais organismos patogênicos podem causar em ostras cultiváveis e prevenir danos à saúde humana e animal.

PALAVRAS-CHAVE

Fungos; bactérias; cultivo de ostras.

RESUMEN

Las ostras representan un alto riesgo para la salud pública porque, al ser organismos filtradores, acumulan grandes cantidades de microorganismos presentes en aguas contaminadas. Por ello, este trabajo tuvo como objetivo investigar la presencia de hongos, levaduras y bacterias aislados de ostras cultivadas mediante una amplia revisión de la literatura publicada durante los últimos diez años. Se realizó una búsqueda bibliográfica en febrero de 2023 utilizando los descriptores: "molusco bivalvo", "cultivo de ostras", "ostricultura", "hongos", "levaduras" y "bacterias". Los estudios seleccionados incluyeron artículos relacionados con la sanidad de las ostras cultivadas en Brasil y en otros países. Se recopiló un total de 719 artículos distribuidos en las bases de datos Science Direct (395), SciELO (06), Google Scholar (127) y PubMed (191). Solo 20 artículos cumplieron con los criterios de inclusión establecidos para la investigación. Los trabajos analizados fueron publicados principalmente en Corea, India y Filipinas durante el año 2020. El estudio revela la escasez de investigaciones relacionadas con hongos y levaduras aislados de ostras cultivadas. Los autores seleccionados identificaron bacterias de los géneros *Vibrio* spp., *Escherichia coli* y *Salmonella* sp. en la mayoría de los estudios. Por lo tanto, es necesario incrementar las investigaciones que permitan evaluar los riesgos que estos organismos patógenos pueden causar en las ostras de cultivo y prevenir daños a la salud humana y animal.

PALABRAS CLAVE

Hongos; bacterias; ostricultura.

INTRODUCTION

Aquaculture is an important source of mollusk production, mainly bivalves. These are species that have a positive environmental impact and bring significant nutritional benefits to consumers (FAO, 2020). The latest edition of the report "The State of World Fisheries and Aquaculture 2024" indicates that global fisheries and aquaculture production in 2022 amounted to 223.2 million t, 4.4% more than in 2020; production included 185.4 million t of aquatic animals and 37.8 million t of algae. Latin America and the Caribbean achieved 17.7 million t of fisheries and aquaculture production, 8% of the world total, rising to 9% when only aquatic animal production is considered (FAO, 2024).

The cultivation of oysters is linked to the environmental conditions of the cultivation area, being subject to the abiotic and biological characteristics (CHAGAS et al., 2021) and the type of management/cultivation (BANNISTER et al., 2019; VALE et al., 2020). Oysters are filter feeders and bioaccumulators, which feed on microalgae, organic and inorganic matter found in the water and retained by the gills, an organ also responsible for absorbing oxygen (SANTOS, BARRETO, SILVA, 2014). Due to the filter feeding habit of oysters, the diet of these animals is based on particles suspended in water, which can cause pathogenic microorganisms present in the production environment to accumulate in the tissues of the bivalve (TROMBETA, NORMANDE, 2017). Therefore, they can accumulate both viruses and bacteria when cultivated in places with the presence of these microorganisms (ABDELZAHER et al., 2009).

The microbiota of oyster meat is directly related to the environment in which it operates. Therefore, these molluscs pose a great risk in their consumption because they are filter feeders and bioaccumulators of microorganisms, which is why they are widely used as bioindicators (SALLES et al., 2017; PHAM et al., 2020). The risks associated with the consumption of molluscs, in many countries, determined the development of a set of own standards in the commercialization process, based on microbiological analyzes of the culture water and/or its meat (SILVA et al., 2004).

According to data from the Ministry of Health's Information and Notifiable Diseases System (SINAN), outbreaks involving fish in Brazil account for 2.1% of cases (SINAN, 2019). The presence of fungi in products of biological origin represents a risk factor for human and animal health, given the potential for producing mycotoxins with carcinogenic, hepatotoxic, neurotoxic and nephrotoxic properties (MONTEIRO, 2018). Thus, oysters are seen as high-risk foods, being largely associated with cases of food poisoning and infection, probably because of increased environmental pollution (NASCIMENTO et al., 2011). The microbiota of most bivalve molluscs is quite diverse and may include viruses and bacteria (Staley et al., 2012). All these pathogens can be transmitted from the environment to humans at the time of ingestion of the mollusk, for this reason, contaminated oysters present a high potential risk to the health of consumers (BRASIL, 2012).

Periodic microbiological analyzes of water from crops and bivalve mollusks make up an excellent parameter indicator of contamination by pathogenic microorganisms. Although the quality of mollusks is dependent on many factors, such as pollution discharge, which can carry toxic substances and various microorganism, toxin -producing microscopic algae and inadequate post-harvest management, water quality is undeniably one of the most important factors (JENKINS et al., 2009).

Several pathogens can be associated with bivalve mollusks causing damage and diseases, with the main organisms belonging to the protozoa group *Sphenophrya* sp., *Trichodina* sp., *Ancistrocoma* sp., *Nematopsis* sp., *Perkinsus* sp., *Tylocephalum* sp. (RLOS) and *Ostracobable implexa* fungus (SCARDUA et al., 2017)

Oysters pose a great risk to consumers, because among marine animals captured in environments contaminated by microorganisms, bivalve molluscs are those that pose the greatest risks to Public Health, as they are filter-feeding and bioaccumulating organisms (FIGUEIREDO et al., 2015). Fungi are dispersed in the environment, in vegetables, atmospheric air, soil, water, in food, in general debris, in animals and in man; mostly, fungi are strict aerobic organisms, except for certain optional anaerobic fermenting yeasts, which can develop in a reduced oxygen environment or even in their absence (MANOHARACHARY et al., 2016).

Given the above, one realizes the importance of a literary basis that assists in the knowledge of the possible organisms acting in the health of oysters. Thus, we aim in this work, conducting research on the presence of fungi, yeast and bacteria isolated from cultivated oysters, through an extensive literature review in the last ten years.

MATERIAL AND METHODS

We conducted a bibliographic survey in the Science Direct databases, periodic portal of the Higher Education Personnel Improvement Coordination (CAPES) and Scientific Electronic Library Online (SciELO), Google Scholar and PubMed in February 2023 using the combination of six Descriptors: "Mollusco Bivalve", "ostriculture", "oyster cultivation", "fungi", "yeast" and "bacteria". The intersection of the descriptors occurred through the boolean operators "or" and "and". The articles published from 2013 to February 2023 were considered. The studies selected to compose this research include articles related to the health of oysters cultivated in Brazil and worldwide. Articles were excluded in which the studies were not conducted from the collection of oysters from cultivation.


After the electronic search procedure in the mentioned databases, the title and summary were read to identify if the articles met the established criteria for inclusion in the research. Then we selected the articles to identify those who composed the final sample of this bibliographic review. Therefore, this study is an exploratory research and, from the point of view of technical procedures, is classified as bibliographic research prepared from material already published (KAUARK, MANHÃES, MEDEIROS, 2010).

RESULTS

Published studies

Overall, we found 719 articles distributed in Science Direct (395), Scielo (06), Google Academic (127) and Pubmed (191) databases. Of the total articles raised, only 20 met the criteria for inclusion in the research. There were no articles that met the inclusion criteria for this research in the Scielo database. Although Asia, with emphasis on Korea, India and the Philippines, concentrated most of the studies in 2020, South America, especially Brazil, also contributed significantly to research in the area, as seen in graph 1. The research focused mainly on the pathogenic bacteria *Vibrio parahaemolyticus* and *Escherichia coli* (graph 2). Only one article reported fungi and yeasts isolated from oysters cultivated in three of the four databases where the research took place. Oysters of the genus *Crassostrea* sp. were the main genus used for the isolation of fungi and bacteria for identification purposes, especially of bacteria of the genus *Vibrio* sp., followed by *Escherichia coli* and *Salmonella* (Table 1 and table 2).

Grafic 1. Continental distribution that the studies were reported.



Grafic 2. Etiological agents reported in the studies found.




Table 1. Articles found in data survey, highlighting the authors (references) database used, research microorganisms, periodicals, and thematic area.

Microorganisms	References	Journal	Thematic area
Science Direct			
Fungi and yeast	SILVA et al., 2020a	Aquaculture Reports	Microbiology
	BRANDÃO et al., 2017	Journal of Food Protection	Microbiology
Bacteria	SILVA et al., 2018b	Marine Pollution Bulletin	Biochemistry
	LORENZONI et al., 2021	Journal of Food Protection	Taxonomy and Ecology
	NEETOO et al., 2022	Journal of Food Protection	Microbiology
	PAKINGKING, 2022	Journal of Food Protection	Microbiology
	NUÑAL et al., 2023	Journal of Food Protection	Microbiology
	OYANEDEL et al., 2023	Journal of Invertebrate Pathology	Taxonomy
Google Scholar			
Bacteria	CHINNADURAI et al., 2016a	Regional Studies in Marine Science	Ecology
	CHINNADURAI et al., 2021b	Aquaculture	Microbiology
	MUDADU et al., 2021	Food Control	Microbiology
PubMed			
Bacteria	NETA et al., 2015	Genetics and Molecular Research	Microbiology
	KANG et al., 2016	Environmental Science and Pollution Research	Ecology
	KIM et al., 2017	Environmental Science and Pollution Research	Ecology
	VEZZULLI et al., 2017	Microbial ecology	Molecular ecology
	AAGESEN et al., 2018	Microbial ecology	Microbiology
	OSTRENSKY et al., 2018	Journal of applied Microbiology	Microbiology
	CANTY et al., 2020	Environmental Microbiology	Taxonomy
	HORODESKY et al., 2020	Anais da Academia Brasileira de Ciências	Taxonomy
	PRUENTE et al., 2020	International Journal of Food Microbiology	Microbiology

Table 2. Summary of the raised studies that are cultivated in the world, ordered by the database.

Microorganisms	References	Host	Etiological agent	Study area
Science Direct				
Fungi and yeast	Silva et al., 2020	<i>Crassostrea gasar</i>	<i>Escherichia coli</i> , <i>positive coagulase staphylococci</i> , mold, yeast, and <i>Salmonella</i> spp.	Pará, Brazil
	BRANDÃO et al., 2017	<i>Crassostrea rhizophorae</i>	<i>Escherichia coli</i> and <i>Vibrio parahaemolyticus</i>	South Coast of Bahia, in the northeast of Brazil
	SILVA et al., 2018	Oyster <i>Crassostrea rhizophorae</i> and mussel <i>Mytella guyanensis</i>	<i>Vibrio parahaemolyticus</i>	Bahia, Brazil
	LORENZONI et al., 2021	Bivalve mollusks <i>Mytilus galloprovincialis</i> , <i>Crassostrea gigas</i> and <i>Ruditapes decussatus</i>	<i>Vibrio parahaemolyticus</i> and <i>Vibrio vulnificus</i>	Sardinia, Italy
Bacteria	NEETOO et al., 2022	Fishes <i>Scarus ghobban</i> , <i>Lethrinus nebulosus</i> , <i>Siganus sutor</i> , <i>Epinephelus fasciatus</i> , <i>Sciaenops ocellatus</i> and <i>Dicentrarchus labrax</i> ; oysters <i>Saccostrea cucullata</i> [It is cultivated]) and sea urchins <i>Tripneustes gratilla</i>	<i>Vibrio alginolyticus</i> , <i>Vibrio parahaemolyticus</i> , <i>Vibrio harveyi</i> and <i>Vibrio diabolicus</i>	Réduit, Moka
	SILVA et al., 2022a	<i>Crassostrea iredalei</i>	<i>Escherichia coli</i> and <i>Salmonella</i>	Province of Capiz, Western Visayas, Philippines
	NUÑAL et al., 2023	Mussel <i>Perna viridis</i> and oysters <i>Margallana bilineata</i>	<i>Escherichia coli</i> , <i>Salmonella</i> spp., <i>Vibrio parahaemolyticus</i> and <i>Vibrio cholerae</i>	Western Visayas, Philippines
	OYANEDEL et al., 2023	<i>Crassostrea gigas</i>	<i>Vibrio</i> sp.	Tongoy Bay, Chile

Google Scholar

Bacteria	CHINNADURAI et al., 2016	<i>Crassostrea madrasensis</i>	Total coliforms, Fecal coliforms, <i>Escherichia coli</i> , <i>Enterococcus faecalis</i>	Ashtamudi Lake (Kerala, India)
	MUDADU et al., 2021	<i>Mytilus galloprovincialis</i> , <i>Crassostrea gigas</i> , <i>Tapes decussatus</i>	<i>Arcobacter</i> spp.	Cultivation Farms in Sardinia (Italy)
	CHINNADURAI et al., 2021b	<i>Crassostrea madrasensis</i>	Total plate counts, Faecal coliforms, Total coliforms, Faecal <i>Streptococci</i> , <i>Escherichia coli</i> , <i>Salmonella</i> spp., and <i>Vibrio</i> spp.	Commercial Farm in Ashtamudi Estuary, Along the Southwest Coast of India

PubMed

Bacteria	NETA et al., 2015	<i>Crassostrea rhizophorae</i>	Coliforms at 35 ° C, Enterobacteriaceae, Aerobic and Psychrotrophic mesophils	Porto do Campo in Camamu Bay - Bahia, Brazil
	KANG et al., 2016	<i>Crassostrea gigas</i>	<i>Vibrio parahaemolyticus</i>	West Coast of Korea
	KIM et al., 2017	<i>Crassostrea gigas</i>	Total and Fecal Coliforms, <i>Escherichia coli</i> <i>Vibrio</i> sp., <i>Pseudoalteromonas</i> , <i>Spirochaetaceae</i> , <i>Alteromonas</i> , <i>Cyanobacteria</i> , <i>Colwellia</i> , <i>Marinomonas</i> , <i>Photobacterium</i> , <i>Klebsiella</i> , <i>Neptunomonas</i> , <i>Rhodobacteraceae</i>	Aquaculture Farm, Korea
	VEZZULLI et al., 2017	<i>Crassostrea gigas</i>		Seafood Farm, Gulf of La Spezia, Italy
	AAGESEN et al., 2018	<i>Crassostrea gigas</i>	<i>Vibrio parahaemolyticus</i>	Oregon Oysters Farm, Newport, Oregon

PubMed

Bacteria	OSTRENSKY et al., 2018	<i>Crassostrea gasar</i>	Bacterial microbiota of the phyllos: Acidobacteria, Actinobacteria, Armatimonadetes, Bacteroidetes, Caldithrix, Chlamydiae, Chlorobi, Chloroflexi, Crenarchaeota, Cyanobacteria, Elusimicrobia, Euryarchaeota, Fibrobacteres, Firmicutes, Fusobacteria, Gemmatimonadetes, Lentisphaerae, Planctomycetes, Proteobacteria, Spirochaetes, Tenericutes, Verrucomicrobia	Marine farms in Guaratuba Bay, on the coast of the state of Paraná, Brazil
	CANTY et al., 2020	<i>Crassostrea virginica</i>	<i>Vibrio vulnificus</i> and <i>Vibrio parahaemolyticus</i>	Bay of Cedar Island, Jarrett Bay, and the Newport River estuary
	HORODESKY et al., 2020	<i>Crassostrea</i> sp.	Bacterial microbiota	Macau and Tibau do Sul in Rio Grande do Norte, Maringá (Paraíba), Passo do Camaragibe and Barra de São Miguel in Alagoas, Brejo Grande and Indiaroba in Sergipe, Brazil
	PRUENTE et al., 2020	<i>Crassostrea virginica</i>	<i>Vibrio vulnificus</i> and <i>Vibrio parahaemolyticus</i>	Cowell, Australia

DISCUSSION

The study reveals the scarcity of research focused on ecology, microbiology, taxonomy or any other aspect involving fungi and yeasts isolated from cultivated oysters. The research conducted by Silva et al. (2020), mentions the isolation of fungi and yeast in *C. Gasar* cultivation in the state of Pará, in Brazil; however, microorganisms were not identified at a specific level, attributing to the taxonomic group co-company as sporadic and unusual in mollusks.

Fungi produce mycotoxins that are secondary metabolic considered harmful to both human and animal health (EISENMAN, CASADEVALL, 2012; IARC, 1993), and some types of these molecules have carcinogenic action and may also interfere with the replication of affected DNA (WHO-IARC, 1993; MAKUN et al., 2011; ISMAIEL, PAPENBROCK, 2015). By ingesting products containing high concentrations of Aflatoxin B1 (AFB1), secondary metabolics produced by some species of fungi of the genus *Aspergillus* sp., the effects may be immunosuppressors, mutagenic, teratogenic, carcinogenic, and hepatotoxic (EISENMAN, CASADEVALL, 2012). Fumonisin, mycotoxins produced by *Fusarium* sp. and ochratoxin A (AO), toxin produced by fungi of *Aspergillus* and *Penicillium* genres are currently classified by IARC (1993) as class 2B, designated to probable carcinogenic agents for humans. Still, shyness is observed when referring to the study of this isolated taxonomic group of cultivated oysters that, in most cases, are consumed in natura. Borzykh and Zvereva, 2018 identified fungi of the genus *Fusarium* sp. and *Penicillium* sp. in oysters *Crassostrea gigas* on Rikord Island, Sea of Japan. The isolation of the genera was also carried out from the scallop *Nodipecten nodosuns* in Angra dos Reis Bay, RJ, Brazil (Santos et al., 2016) and in scallop *Mizuhopecten yessoensis* in Rikord Island, Sea of Japan (Borzykh and Zvereva, 2018). The genus *Penicillium* was also found in gill arches, intestine and muscle of the scallop *Nodipecten nodosuns* in Brazil (Santos et al., 2016), in the mantle, adductor muscle, kidneys, gills, hepatopancreas, and gonads of mussels *Modiolus modiolus* and *Crenomytilus grayanus* in Ussuriiskii Bay, Peter the Great Bay, Sea of Japan (Zvereva and Vysotskaya, 2005), scallop *Mizuhopecten yessoensis* and mussel *Mytilus trossulus* in Rikord Island, Sea of Japan (Borzykh and Zvereva, 2018). The genus *Aspergillus* has already been identified in Brazil isolated from other mollusks such as scallop *Nodipecten nodosuns* (Santos et al., 2016), in Japan in scallop *Mizuhopecten yessoensis* (Borzykh and Zvereva, 2018) and in mussels *Modiolus modiolus* and *Crenomytilus grayanus* also in Japan (Zvereva and Vysotskaya, 2005).

Bacteria of the genus *Vibrio* spp. were identified in most of the work. Probably this factor is associated with the epidemiological importance of the genre (AUSTIN, 2010), as some species are considered pathogenic for human health such as *V. parahaemolyticus* and *V. vulnifus* (AUSTIN, 2010; SOUSA, 2004; SILVA et al., 2018). These organisms were the most reported in the studies tabulated in the present research. Both species cause infections commonly related to ingestion of raw or poorly cooked animals (OLIVER, 2013).

The bacterium *Vibrio parahaemolyticus* was isolated in 45% of studies, being detected in a variety of marine organisms and geographic regions. Among the hosts, fish (such as *Scarus ghobban*, *Lethrinus nebulosus*, *Siganus sutor*, *Epinephelus fasciatus*, *Sciaenops ocellatus* and *Dicentrarchus labrax*), bivalve molluscs (oysters of the genera *Saccostrea*, *Crassostrea* and *Magallana*, mussels of the genera *Mytella*, *Perna* and *Mytilus*) and sea urchins (*Triplaneustes gratilla*) stand out. The presence of *V. parahaemolyticus* has been confirmed in different regions, including the Indian Ocean (Réduit, Moka), the Atlantic Ocean (Southern coast of Bahia, Brazil; Cedar Island Bay, Jarrett Bay and Newport River estuary, United States; West coast of Korea; Oregon Oyster Farm, Newport, Oregon; Cowell, Australia) and the Pacific Ocean (Western Visayas, Philippines; Sea of Japan; Sardinia, Italy), as shown in Table 2. *V. parahaemolyticus* is considered a zoonosis with more frequent cases of infections, although lighter; in humans can generate diarrhea with abdominal cramps, nausea, vomiting and fever (DRAKE., DEPAOLA, JAYKUS, 2007; JONES, OLIVER, 2009), besides to find marine animals (VASEEHRAN, RAMASAMY, 2003). *V. vulnifus* cases are considered less frequent, however, their effects may be more severe, especially in immunocompromised individuals, and as symptoms of infection can cause fever, hypotension, and chills (YEUNG, BOOR, 2004), while *Salmonella* sp. and *E. coli*, if found in animals, are indicative of fecal contamination (LEES, 2000), but all species are constantly associated with food transmitted diseases that can raise risks to human health (LEES, 2000; OLIVEIRA et al, 2011). The results obtained in this study

reveal a high prevalence of pathogenic bacteria in oysters and reinforce the importance of continued research to understand the ecology and epidemiology of these bacteria, as well as to develop effective strategies for the control and prevention of foodborne diseases.

The constant frequency of studies with oysters of the genus *Crassostrea* sp. It is directly related to its source of nutrients (SOLETCHNIK et al., 2002) and ease of cultivation (FAO, 2014), which guarantee their broad marketing worldwide (FAO, 2014). These factors can be unfeasible when oysters become susceptible to the presence of pathogens when cultivated, and this reflects the need for increased studies and consequently measures that may minimize the risks to human health in the consumption of products originating from aquaculture (CHINNADURAI et al., 2021).

CONCLUSION

The study reveals that there is a scarcity of research aimed at identifying fungi and isolated yeasts from cultivated oysters, even considering the aggravation that these microorganisms are producing toxins potentially carcinogenic to man, especially because oysters are consumed mainly in raw form.

Bacteria of the genres *Vibrio* spp. and *Salmonella* sp., and the *E. coli* species are commonly associated with cultivation oysters and are configured in the main pathogenic organisms that cause concern for public health.

Ostreculture is growing significantly and is accompanied by the need to obtain research that aims at the broader knowledge of the health of these organisms, especially because oysters have bioaccumulation potential, making them an important reservoir of pathogens.

PATENTS

Funding: This research received no external funding.

Acknowledgments: The authors would like to thank the Laboratory Microbiology, parasitology, and biotechnology (MIPABIO) of State University of Maranhão (UEMA).

Conflicts of Interest: "The authors declare no conflict of interest."

REFERENCES

- AAGESEN, A. M.; PHUVASATE, S.; SU, Y. C.; HÄSE, C. C. Characterizing the adherence profiles of virulent *Vibrio parahaemolyticus* isolates. **Microbial ecology**, v. 75, p. 152-162. 2018. DOI: 10.1007/s00248-017-1025-8
- ABDELZAHER, A. M.; WRIGHT M. E.; ORTEGA C.; SOLO-GABRIELE H. M.; MILLER G.; ELMIR S.; et al. Presence of pathogens and indicator microbes at a non-point source subtropical recreational marine beach. **Applied and environmental microbiology**, v. 76, n. 3, p. 724-732. 2010. DOI: 10.1128/aem.02127-09
- AUSTIN, B. Vibrios as causal agents of zoonoses. *Veterinary microbiology*, v. 140, n. 3-4, p. 310-317. 2010. DOI: 10.1016/j.vetmic.2009.03.015
- BANNISTER J.; SIEVERS M.; BUSH F.; BLOECHER N. Biofouling in marine aquaculture: a review of recent research and developments. *Biofouling*, v. 35, n. 6, p. 631-648. 2019. DOI: 10.1080/08927014.2019.1640214
- BRANDÃO M. A. R.; LOPES A. I. S. T.; NETA M. R. S. T.; OLIVEIRA R. B. F.; REZENDE R. P.; ALBUQUERQUE G. R.; GONÇALVES, V. D.; RODRIGUES, D. D. P.; BOEHS, G.; MACIEL, B. M. Microbiological quality and prevalence of-lactam antibiotic resistance genes in oysters (*Crassostrea rhizophorae*). *Journal of Food Protection*, v. 80, n. 3, p. 488-496. 2017. DOI: 10.4315/0362-028X.JFP-16-098

BRASIL – Ministério da Pesca e Aquicultura. Ministério da Agricultura, Pecuária e Abastecimento. Instrução Normativa Interministerial MPA/MAPA N° 7, de 08 de maio de 2012. In: Diário Oficial da União, n. 89, seção 1, p. 55-59, 09 de maio de 2012.

CANTY, R.; BLACKWOOD, D.; NOBLE, R.; FROELICH, B. A comparison between farmed oysters using floating cages and oysters grown on bottom reveals more potentially human pathogenic *Vibrio* in the on bottom oysters. *Environmental microbiology*, v. 22, n. 10, p. 4257-4263. 2020. DOI: 10.1111/1462-2920.14948

CHAGAS R. A.; BARROS M. R. F.; ABREU V. S.; SILVA R. E. O.; ALMEILDA M. S.; SANTOS W. C. R.; HERRMANN, M. Growth Performance of the Mangrove Oyster Cultivated on the Amazon Coast. *Brazilian Archives of Biology and Technology*, v. 64.2021. DOI: 10.1590/1678-4324-2021190650

CHINNADURAI, S.; ELAVARASAN, K.; GEETHALAKSHMI, V.; KRIPA, V.; MOHAMED, K. S. Evaluation of static and flow-through depuration system on depuration of naturally contaminated farmed edible oyster *Crassostrea madrasensis* (Preston, 1916). *Aquaculture*, v. 545, p. 737141. 2021. DOI: 10.1016/j.aquaculture.2021.73714

CHINNADURAI, S.; MOHAMED, K. S.; SHARMA, J.; VENKATESAN, V.; KRIPA, V. Assessment of bio-accumulation of bacteria in oysters from shellfish growing waters in Ashtamudi Lake (Kerala, India): A RAMSAR wetland. *Regional Studies in Marine Science*, v. 7, p. 118-122. 2016. DOI: 10.1016/j.rsma.2016.05.016

DRAKE, S. L.; DEPAOLA, A.; JAYKUS, Lee-Ann. An overview of *Vibrio vulnificus* and *Vibrio parahaemolyticus*. *Comprehensive reviews in food science and food safety*, v. 6, n. 4, p. 120-144. 2007. DOI: 10.1111/j.1541-4337.2007.00022.x

EISENMAN, H. C.; CASADEVALL, A. Synthesis and assembly of fungal melanin. *Applied microbiology and biotechnology*, v. 93, p. 931-940. 2012. DOI: 10.1007/s00253-011-3777-2

FAO - Food And Agriculture Organization Of The United Nations. The state of world fisheries and aquaculture: opportunities and challenges. Roma: FAO, 2014.

FAO - Food And Agriculture Organization. The state of world fisheries and aquaculture. Rome: FAO. p. 227, 2020.

FAO - Food And Agriculture Organization. O estado da pesca e da aquicultura no mundo 2024. Roma: FAO, 2024.

FIGUEIREDO, J. F.; RIBEIRO, S.; PAULA, M; PONTES, A. Determinação da concentração de Coliformes Totais e Termotolerantes na água de Cultivo de ostras do mangue (*Crassostrea rhizophorae*) em região estuarina. *Enciclopédia Biosfera, Centro Científico Conhecer*, v.11, n.21, p. 3498, 2015.

HORODESKY, A.; CASTILHO-WESTPHAL, G. G.; PONT, G. D.; FAORO, H.; BALSANELLI, E.; TADRA-SFEIR, M. Z.; COZER, N.; PIE, M. R.; OSTRENSKY, A. Metagenomic analysis of the bacterial microbiota associated with cultured oysters (*Crassostrea* sp.) in estuarine environments. *Anais da Academia Brasileira de Ciências*, v. 92. 2020. DOI: 10.1590/0001-3765202020180432

IARC World Health Organization International Agency For Research On Cancer. *Toxins derived from Fusarium moniliforme: fumonisins B1 and B2 and fusarin C*. *IARC Monographs on the Evaluation of Carcinogenic Risks to Humans*. v. 56, 1993. p. 445-462.

ISMAIEL, A. A.; PAPENBROCK, J. Mycotoxins: producing fungi and mechanisms of phytotoxicity. *Agriculture*, v. 5, n. 3, p. 492-537. 2015. DOI: 10.3390/agriculture5030492

JENKINS, M. W.; TIWARI, S.; LORENTE, M.; GICHABA, C. M.; WUERTZ, S. Identifying human and livestock sources of fecal contamination in Kenya with host-specific Bacteroidales assays. *Water Research*, v. 43, n. 19, p. 4956-4966. 2009. DOI: 10.1016/j.watres.2009.07.028

JONES M. K.; OLIVER J. D. *Vibrio vulnificus*: disease and pathogenesis. *Infection and immunity*, v. 77, n. 5, p. 1723-1733. DOI: 10.1128/IAI.01046-08

KANG, C. H.; SHIN, Y.; KIM, W.; KIM, Y.; SONG, K.; OH, E. G.; KIM, S.; YU, H.; SO, J. S. Prevalence and antimicrobial susceptibility of *Vibrio parahaemolyticus* isolated from oysters in Korea. *Environmental Science and Pollution Research*, v. 23, p. 918-926. 2016. DOI: 10.1007/s11356-015-5650-9

KAUARK, F. D. S.; MANHÃES, F. C.; MEDEIROS, C. H. Metodologia da pesquisa: um guia prático. 2010. p. 89.

KIM, J. H.; SHIM, K. B.; SHIN, S. B.; PARK, K.; OH, E. G.; SON, K. T.; YU, H.; LEE, H. J.; MOK, J. S. Comparison of bioaccumulation and elimination of *Escherichia coli* and male-specific bacteriophages by ascidians and bivalves. *Environmental Science and Pollution Research*, v. 24, p. 28268-28276. 2017. DOI: 10.1007/s11356-017-0736-1

LEES, D. Viruses and bivalve shellfish. *International journal of food microbiology*, v. 59, n. 1-2, p. 81-116. 2000.

LORENZONI, G.; TEDDE, G.; MARA, L.; BAZZONI, A. M.; ESPOSITO, G.; SALZA, S.; PIRAS, G.; TEDDE, T.; BAZZARDI, R.; ARRAS, I.; UDA, M., T.; VIRGILIO, S.; MELONI, D.; MUDADU, A. G. Presence, seasonal distribution, and biomolecular characterization of *Vibrio parahaemolyticus* and *Vibrio vulnificus* in shellfish harvested and marketed in Sardinia (Italy) between 2017 and 2018. **Journal of Food Protection**, v. 84, n. 9, p. 1549-1554, 2021. DOI: 10.4315/JFP-21-059

MAKUN, H. A.; DUTTON, M. F.; NJOBEB, P. B.; MWANZA, M.; KABIRU, A. Y. Natural multi-occurrence of mycotoxins in rice from Niger State, Nigeria. *Mycotoxin research*, v. 27, p. 97-104. 2011. DOI: 10.1007/s12550-010-0080-5

MANOHARACHARY C.; TILAK K. V. B. R.; MALLAIAH K. V.; KUNWAR I. K. *Mycology and microbiology (A textbook for UG and PG courses)*. Scientific Publishers, 2016. p. 606.

MONTEIRO, J. P. S.; BOSSA, L. F.; MENDONÇA, M. F. F. F.; BALDO, C.; FUNGARO, M. H. P.; SARTORI, D. Genetic diversity and macromorphological variability of *Aspergillus* isolates from garlic collected in different brazilian states. In: ANAIS DO III CONGRESSO PARANAENSE DE MICROBIOLOGIA, 2018, Londrina. Anais eletrônicos... Campinas, Galoá, 2018. Disponível em: <<https://proceedings.science/acpm/cpm-2018/trabalhos/genetic-diversity-and-macromorphological-variability-of-aspergillus-isolates-fro?lang=pt-br>> Acesso em: 17 Set. 2024.

MUDADU, A. G.; SALZA, S.; MELILLO, R.; MARA, L.; PIRAS, G.; SPANU, C.; TERROSU, G.; FADDA, A.; VIRGILIO, S.; TEDDE, T. Prevalence and pathogenic potential of *Arcobacter* spp. isolated from edible bivalve molluscs in Sardinia. *Food Control*, v. 127, p. 108139. 2021. DOI: 10.1016/j.foodcont.2021.108139

NASCIMENTO V. A.; MITTARAQUIS A. S. P.; TRAVÁLIA B. M.; SANTOS R. C. A.; NUNES M. L.; DE AQUINO L. C. L. Qualidade Microbiológica de Moluscos Bivalves-Sururu e Ostras submetidos a tratamento térmico e estocagem congelada. *Scientia Plena*, v. 7, n. 4. 2011.

NEETOO H.; REEGA K.; MANOGA Z. S.; NAZURALLY N.; BHOYROO V.; ALLAM M.; et al. Prevalence, genomic characterization, and risk assessment of human pathogenic *Vibrio* Species in Seafood. *Journal of Food Protection*, v. 85, n. 11, p. 1553-1565. 2022. DOI: 10.4315/JFP-22-064

NETA, M. S.; MACIEL, B. M.; LOPES, A. T. S.; MARQUES, E. L. S.; REZENDE, R. P.; BOEHS, G. Microbiological quality and bacterial diversity of the tropical oyster *Crassostrea rhizophorae* in a monitored farming system and from natural stocks. *Genetics and Molecular Research*, v. 14, n. 4, p. 15754-15768. 2015. DOI: 10.4238/2015.December.1.27

NUÑAL S. N.; MONAYA K. J. M.; MUEDA C. R. T.; SANTANDER-DE LEON S. M. Microbiological Quality of Oysters and Mussels Along Its Market Supply Chain. *Journal of Food Protection*, v. 86, n. 3, p. 100063. 2023. DOI: 10.1016/j.jfp.2023.100063

OLIVEIRA, J.; CUNHA, A.; CASTILHO, F.; ROMALDE, J. L.; PEREIRA, M. J. Microbial contamination and purification of bivalve shellfish: Crucial aspects in monitoring and future perspectives—A mini-review. *Food Control*, v. 22, n. 6, p. 805-816. 2011. DOI: 10.1016/j.foodcont.2010.11.032

OLIVER, J. D. *Vibrio vulnificus*: death on the half shell. A personal journey with the pathogen and its ecology. *Microbial ecology*, v. 65, p. 793-799. 2013. DOI: 10.1007/s00248-012-0140-9

OSTRENSKY, A.; HORODESKY, A.; FAORO, H.; BALSANELLI, E.; SFEIR, M. Z. T.; COZER, N.; PIE, M. R.; DAL PONTE, G.; CASTILHO-WESTPHAL, G. G. Metagenomic evaluation of the effects of storage conditions on the bacterial microbiota of oysters *Crassostrea gasar* (Adanson, 1757). *Journal of applied microbiology*, v. 125, n. 5, p. 1435-1443. 2018. DOI: 10.1111/jam.14045

OYANEDEL D.; ROJAS R.; BROKORDT K.; SCHMITT P. *Crassostrea gigas* oysters from a non-intensive farming area naturally harbor potentially pathogenic *vibrio strains*. *Journal of Invertebrate Pathology*, v. 196, p. 107856. 2023. DOI: 10.1016/j.jip.2022.107856

PAKINGKING JR R.; HUALDE M. L.; PERALTA E.; FAISAN J.; USERO R. Microbiological quality and heavy metal concentrations in slipper oyster (*Crassostrea iredalei*) cultured in major growing areas in Capiz Province, Western Visayas, Philippines: Compliance with international shellfish safety and sanitation standards. *Journal of Food Protection*, v. 85, n. 1, p. 13-21. 2022. DOI: 10.4315/JFP-21-257

PHAM, L. T.; HOANG, T. T. T.; TU, L. C. T.; TRAN, Y. H. T.; LE, B. D.; VAN NGUYEN; XUAN DO, H; VAN THAI, N. Bioaccumulation and health risk assessment of polycyclic aromatic hydrocarbons in oyster (*Crassostrea* sp.) and gastropod (*Cymatium* sp.) species from the Can Gio Coastal Wetland in Vietnam. *Marine and Freshwater Research*, v. 71, n. 6, p. 617-626, 2019. DOI: 10.1071/MF19055

PRUENTE, V. L.; JONES, J. L.; STEURY, T. D.; WALTON, W. C. Effects of tumbling, refrigeration and subsequent resubmersion on the abundance of *Vibrio vulnificus* and *Vibrio parahaemolyticus* in cultured oysters (*Crassostrea virginica*). *International Journal of Food Microbiology*, v. 335, p. 108858. 2020. DOI: 10.1016/j.ijfoodmicro.2020.108858

SABRY, R. C.; MAGALHÃES, A. R. M. Parasitas em ostras de cultivo (*Crassostrea rhizophorae* e *Crassostrea gigas*) da Ponta do Sambaqui, Florianópolis, SC. *Arquivo Brasileiro de Medicina Veterinária e Zootecnia*, v. 57, p. 194-203. 2005. DOI: 10.1590/S0102-09352005000800010

SANTOS, S. S. dos; BARRETO, N. S. E.; SILVA. Delineamento do perfil dos consumidores e classificação dos estabelecimentos que comercializam ostras em Valença, Bahia, Brasil. *Arquivos de Ciências do Mar*, v.2, n. 47, p. 72-79. 2014.

SCARDUA M. P.; VIANNA R. T.; DUARTE S. S.; FARIAS N. D.; CORREIA M. L. D.; SANTOS H. T. A. D.; SILVA P. M. D. Growth, mortality and susceptibility of oyster *Crassostrea* spp. to *Perkinsus* spp. infection during on growing in northeast Brazil. *Revista Brasileira de Parasitologia Veterinária*, v. 26, p. 401-410. 2017. DOI: 10.1590/S1984-29612017061

SALLES, P.B.D.; MACEDO, Y.B.; FIGUEIREDO, E.L. Caracterização físicoquímica e microbiológica da carne do molusco Bivalve Sarnambi (*Phacoides pectinitus*) coletado nas praias em Algodual e Salinópolis, no Pará. *Revista Brasileira de Tecnologia Agroindustrial*, v. 11, n. 1, p. 2245-2261, 2017. DOI: 10.3895/rbta.v11n1.2907

SIEVERS M.; DEMPSTER T.; KEOUGH M. J, FITRIDGE I. Methods to prevent and treat biofouling in shellfish aquaculture. *Aquaculture*, v. 505, p. 263-270. 2019. DOI: 10.1016/j.aquaculture.2019.02.071

SILVA A. I.; VIEIRA R. H.; MENEZES F. G.; FONTELES-FILHO A. A.; TORRES R. C.; SANT'ANNA E. S. Bacteria of fecal origin in mangrove oysters (*Crassostrea rhizophorae*) in the Cocó River estuary, Ceará State, Brazil. *Brazilian Journal of Microbiology*, v. 35, p. 126-130. 2004. DOI: 10.1590/s1517-83822004000100021

SILVA I. P.; CARNEIRO C. S.; SARAIVA M. A. F.; OLIVEIRA T. A. S.; SOUSA O. V.; EVANGELISTA-BARRETO N. S. Antimicrobial resistance and potential virulence of *Vibrio parahaemolyticus* isolated from water and bivalve mollusks from Bahia, Brazil. *Marine Pollution Bulletin*, v. 131, p. 757-762. 2018. DOI: 10.1016/j.marpolbul.2018.05.007

SILVA O. L. L.; VERÍSSIMO S. M. M.; ROSA A. M. B. P.; IGUCHI Y. B.; NUNES E. S. C. L. N.; MORAES C. M., et al. Effect of environmental factors on microbiological quality of oyster farming in Amazon estuaries. *Aquaculture Reports*, v. 18, p. 100437. 2020. DOI: 10.1016/j.aqrep.2020.100437

SOLECHNIK, P.; HUVET, A.; LE MOINE, O.; RAZET, D.; GEAIRON, P.; FAURY, N.; GOULLETQUER, P.; BOUDRY, P. A comparative field study of growth, survival and reproduction of *Crassostrea gigas*, *C. angulata* and their hybrids. *Aquatic Living Resources*, v. 15, n. 4, p. 243-250. 2002. DOI: 10.1016/S0990-7440(02)01175-0

SOUSA O. V.; VIEIRA R. H. S. F.; MENEZES F. G. R.; REIS C. M. F.; HOFER E. Detection of *Vibrio parahaemolyticus* and *Vibrio cholerae* in oyster, *Crassostrea rhizophorae*, collected from a natural nursery in the Cocó river estuary, Fortaleza, Ceará, Brazil. *Revista do Instituto de Medicina Tropical de São Paulo*.v. 46, p. 59-62. 2004. DOI: 10.1590/S0036-46652004000200001

STALEY, C.; RECKHOW, K. H.; LUKASIK, J.; HARWOOD, V. J. Assessment of sources of human pathogens and fecal contamination in a Florida freshwater lake. *Water Research*, v. 46, n. 17, p. 5799-5812, 2012.

TRAVERS M. A.; MILLER K. B.; ROQUE A.; FRIEDMAN C. S. Bacterial diseases in marine bivalves. *Journal of invertebrate pathology*, v. 131, p. 11-31. 2015. DOI: 10.1016/j.jip.2015.07.010

TROMBETA, T. D.; NORMANDE, A. C. L. Microbiological evaluation of grown on the coast of Alagoas State submitted to purification in a closed recirculation system. *Acta of Fisheries and Aquatic Resources*, v. 5, n. 3, p. 48-53. 2017. DOI: 10.2312/ActaFish.2017.5.3.48-53

VALE A. V. P.; SANTOS W. C. R.; BARROS M. R. F.; CHAGAS R. A.; HERRMANN M. Comparação de substratos artificiais na redução de bioincrustantes em um cultivo de ostras no estuário amazônico. *Revista CEPSUL-Biodiversidade e Conservação Marinha*, v. 9, p. e2020001-e2020001. 2020. DOI: 10.37002/revistacepsul.vol9.875e2020001

VASEEHARAN, B. A. R. P.; RAMASAMY, P. Control of pathogenic *Vibrio* spp. by *Bacillus subtilis* BT23, a possible probiotic treatment for black tiger shrimp *Penaeus monodon*. *Letters in applied microbiology*, v. 36, n. 2, p. 83-87. 2003. DOI: 10.1046/j.1472-765X.2003.01255.x

VASQUEZ G. A.; DE SA S. H. G.; SILVA G. S.; BALLESTEROS J. E. M.; BARBIERI E.; SOUSA R. L. M.; Qualidade microbiológica de moluscos e avaliação de EC seca compacta para detecção de coliformes totais e *Escherichia coli*. *Acta Alimentos*. v. 49, p. 32-39. 2020. DOI: 10.1556/066.2020.49.1.5

VEZZULLI, L.; STAGNARO, L.; GRANDE, C.; TASSISTRO, G.; CANESI, L.; PRUZZO, C. Comparative 16SrDNA gene-based microbiota profiles of the Pacific oyster (*Crassostrea gigas*) and the Mediterranean mussel (*Mytilus galloprovincialis*) from a shellfish farm (Ligurian Sea, Italy). *Microbial ecology*, v. 75, p. 495-504. 2018. DOI: 10.1007/s00248-017-1051-6

WESTPHAL, G. G. C.; OSTRENSKY, A. Use of side-scan sonar for estimations of *Crassostrea brasiliiana* (Lamarck, 1819) stocks in subtidal banks on the south coast of Brazil. *Brazilian Journal of Oceanography*, v. 64, p. 49-56. 2016. DOI:10.1590/S1679-87592016104506401

YEUNG, P. M.; BOOR, K. J. Epidemiology, pathogenesis, and prevention of foodborne *Vibrio parahaemolyticus* infections. *Foodborne Pathogens & Disease*, v. 1, n. 2, p. 74-88. 2004. DOI: 10.1089/153531404323143594

-
1. Postgraduate Program in Legal Amazon Biodiversity and Biotechnology, State University of Maranhão, São Luís, Maranhão, Brazil; <https://orcid.org/0000-0003-2858-4549>
 2. Postgraduate Program in Legal Amazon Biodiversity and Biotechnology, State University of Maranhão, São Luís, Maranhão, Brazil; <https://orcid.org/0000-0001-6309-8787>
 3. Program in Ecology and Biodiversity Conservation, São Luís, Maranhão, Brazil; <https://orcid.org/0000-0002-0069-7253>
 4. Department of Biology, State University of Maranhão, São Luís, Maranhão, Brazil. <https://orcid.org/0000-0003-0796-6388>
 5. Program in Ecology and Biodiversity Conservation, São Luís, Maranhão, Brazil; <https://orcid.org/0000-0003-1622-5434>
 6. Postgraduate Program in Legal Amazon Biodiversity and Biotechnology, State University of Maranhão, São Luís, Maranhão, Brazil; <https://orcid.org/0000-0001-9055-1911>
-

Recebido em: 14 de Fevereiro de 2024

Avaliado em: 29 de Outubro de 2025

Aceito em: 10 de Janeiro de 2026



www.periodicos.uniftc.edu.br



Periódico licenciado com Creative Commons
Atribuição-NãoComercial 4.0 Internacional.